



MAKATS E.F.

VEGETATIVE DIAGNOSTICS ACCORDING TO V. MAKATS. BIOPHYSICAL PECULIARITIES

PhD, associate professor (Vinnytsia National medical university named after Pirogov M.I., Ukraine)

These materials are unparalleled and is the intellectual property of developers towards "functional-vegetative diagnosis without using external power sources" (VG Makaz, Makaz EF, Makaz Dm. V., Makaz Dan. C.). Considered its biophysical characteristics and objectivity in assessing the effectiveness of rehabilitation.

Keywords: Acupuncture therapy, functional diagnostics autonomic, vegetative homeostasis, functional rehabilitation.

Наведені матеріали не мають аналогів і є інтелектуальною власністю розробників напрямку "Функціонально-вегетативна діагностика без використання зовнішніх джерел струму" (Макац В. Г., Макац Є. Ф., Макац Дм. В., Макац Ден. В.). Розглянуті її біофізичні особливості і об'єктивність при оцінці реабілітаційної ефективності.

Ключевые слова: Чжень-цзю терапия, функционально-вегетативная диагностика, вегетативный гомеостаз, функциональная реабилитация.

Приведенные материалы не имеют аналогов и является интеллектуальной собственностью разработчиков направления "Функционально-вегетативная диагностика без использования внешних источников тока" (Макац В. Г., Макац Е. Ф., Макац Дм. В., Макац Дэн. В.). Рассмотрены ее биофизические особенности и объективность при оценке реабилитационной эффективности.

Ключевые слова: Чжень-цзю терапия, функционально-вегетативная диагностика, вегетативный гомеостаз, функциональная реабилитация.

Introduction

Before analyzing these histograms remind the international nomenclature acupuncture channels (MAN, Figure 1), which we always enjoy. In this case, we note that submissions are unparalleled [1-22].

A methodology of functional-vegetative diagnosis poses a real control of the therapeutic and rehabilitation effectiveness.

Before studying functional-vegetative system of human, we need to get acquainted with the methodology, which allowed discovering previously unknown biophysical reality...

Till the recent days, vegetative anamnesis with the usage of questionnaire tables has been generally accepted as the means of integral evaluation of vegetative homeostasis. At the same time, historically preserved detached evaluation of sympathetic and parasympathetic reactions of the basic divisions of vegetative nervous system (VNS) contradicts contemporary ideas of their functional-dependent composition. Additionally, complexity of study of super-segmental and segmental levels of vegetative regulation restricts necessary information for clinicians. It is necessary to remember about heterogeneity of sym-

Traditional channel	IAN *	FN	Traditional channel	IAN *	FN
Lungs	LU	P	Urinary bladder	BL	V
Large intestine	LI	GI	Kidney	KI	R
Stomach	ST	E	Pericardium	PC	MC
Spleen – Pancreas	SP	RP	Triple energizer	TE	TR
Heart	HT	C	Gall bladder	GB	VB
Small intestine	SI	IG	Liver	LR	F

*IAN - International Acupuncture Nomenclature (WHO); FN - its French analogue.

Fig 1 Abbreviation of acupunctural channels

pathetic and parasympathetic reactions, nonlinearity of parameters of vegetative status during the change of activity of one of the divisions of VNS and their dependency on a range of regulating factors. At the same time, it has been defined (Wane 2000; Nozdrahov, 2003), that the indications of instrumental examination of VNS characterize only separate mechanisms of vegetative regulation.

Thus, table methods of vegetative diagnostics are subjective, and instrumental reflect functional state of separate subsystems of VNS and separate mechanisms of vegetative regulations. That is why, in order to see the full picture we need enormous, simultaneous and complex researches, during which, separate indications lose features of system evaluation of general vegetative homeostasis (VH).

Our elaborated methodology of functional-vegetative diagnostics (FVD) allowed identifying acupunctural channels, and discovers previously unknown functional-vegetative system of human. For today, this is the only “electropunctural” diagnostics, results of which are stable and comparable in time. It is grounded on the previously unknown biophysical phenomena, with original standardization and directed at evaluation of functional-vegetative homeostasis: correlation of syndromes of sympathetic (*YANG*) and parasympathetic (*YIN*) activity.

We start with the statement, that every electropunctural diagnostics has the right to exist, at least under three conditions:

- 1) presence of testing (diagnostic) signal, power-informational characteristics of which do not exceed biophysical parameters of the system under attention;
- 2) concrete understanding of the functional specifics of the subject and the area of its biophysical influence;
- 3) availability of comparable results of functional diagnostics during repeated (in 5-10-20 min.) examination.

At the same time, we should pay attention to significant mistake of the officially recognized electropunctural technologies, that “condition diagnoses” of a separate organ or separate functional system!


In our case the talk is about dynamic stability of intersystem dependency, i.e. about functional-vegetative homeostasis, forms and measures of its functional deflections. Any other interpretation of received information from representative acupunctural zones leads us from the Eastern metaphysics to the Western one. In other words, contemporary interpretation of the consequences of electropunctural diagnostics is biophysically incorrect.

And the last one. It is unreasonable and useless to argue with the developers of various electropunctural modifications (but, it should be admitted, that discovered biophysical realities, today, cannot be correctly described in the frames of classical biophysics and clinical physiology). Anyway, any opponent can individually check and challenge any technology by testing its comparability of repeated results... In such cases, our experience tells us, that it is the best way to omit verbal battles.

The methodology of functional-vegetative diagnostics (FVD) is described in details in the monographs [1-5], that is why we are going observe only its technical and biophysical peculiarities.

Technical peculiarities of functional-vegetative diagnostics

Diagnostic complexes VITA-01-M and computerized system VITA-01-Biotest (fig. 1) are recommended for usage in practical medicine RPC HM of Ukraine “New medical technology and new means for diagnostics, prevention and rehabilitation” (record №5, 25.12.91) and Scientific council of HM of Ukraine (record №1.08-01, 11.01. 94).



System VITA-01-M

Complex VITA-01-BIOTEST

Fig.1 Equipment for FVD According to V. Makats

Reasonability of functional-vegetative examination of children is confirmed by the program “Two-stage system of rehabilitation of vegetative disorders of children, that live in the zone of ecological control of Ukraine” (executed according to the Assignment of Cabinet of Ministers of Ukraine №1861/4, 4.04.1997 and №12010/87, 01.06. 1999).

Systems VITA-01-M do not require metrological standardization because:

- 1) methodology FVD does not require external sources of power;
- 2) voltage of closed individually-diagnostic circuit does not exceed the level of membrane potential (0,03-0,6 V);
- 3) analysis is needed not for absolute values of diagnostic indices, but for relative correlation of total activity of functional systems *YANG/YIN* groups (syndromes of sympathetic / parasympathetic activity).

Factor of attention of FVD is ability of biological system to generate current into external closed circuit “electrode-donor of electrons (**DE**) – biological object – electrode acceptor of electrons (**AE**)”. At the same time, it should be remembered, that electromagnetic factors of external sources significantly exceed biophysical level of cellular membranes and condition expected excitation (oppression) of acupunctural zones.

It is incorrect to take the reason as the basis to talk about the functional probability of results. Additionally, biodynamics of every acupunctural zone has individual frequency profile, which by mistake is interpreted from “diagnostic” point of view.

Standardization framework of FVD is specific. It is related to evaluation of functional-vegetative equilibrium and is conditioned by statistically probable amount of observations (14. 304 examined children of different age and gender groups).

Methodological peculiarities of functional vegetative diagnostics

Methodological peculiarities of FVD are conditioned by:

- a) short (3 sec.) contact of paired diagnostic electrode DE with symmetric representative zones and limitation of number of tests from 24 to 12;
- b) wet electrode contact with acupunctural zones (neutralizes vegetative-vascular reactions of skin);
- c) usage for electrode AE of the centralized “basic zone” (navel region, equidistant from the zones of representative contact).

At the same time, attention of FVD is concentrated on bioelectrical activity of symmetrical zones-accomplices (tai-yuan, da-ling, shen-men, wan-gu, yang-chi, yang-xi,

tai-bai, tai-chong, tai-xi, shu-gu, qiu-xu and chong-yang). Their individual resistance to direct current is equivalent to average resistance of other single-channel zones (J.Nakatani).

The received in mV (mkA) data of FVD is transformed into average values. We determine overall bioelectric activity of functional systems *YANG* and *YIN* groups, and vegetative coefficient of their interdependency ($k = \sum Y\text{ANG} : \sum Y\text{IN}$). From the point of view of vegetative homeostasis, the latter points to correlation of sympathetic (*YANG*) and parasympathetic (*YIN*) functional activity.

Functional-vegetative principles of FVD. From biophysical point of view, functional activity of separate acupunctural zones is not a carrier of basic information. However, interrelation *YANG/YIN* syndromes directly shows the prevalence of systemic excitation (sympathetic orientation of functional-vegetative activity), or oppression (parasympathetic orientation of functional-vegetative activity).

This conditions the following principles of FVD):

- overall bioelectrical activity of representative zones of group *YANG* is comparable with sympathetic activity, and of group *YIN* – with parasympathetic;

- sympathetic and parasympathetic divisions of VNS at organ level ensure biochemical control over systemic vegetative equilibrium (the latter under ordinary conditions is dynamically stable; disorder of equilibrium is conditioned by prevalence of activity of one of divisions of VNS);

- correlation of *YANG* and *YIN* syndromes at biophysical level characterizes functional vegetative equilibrium, i. e. interdependency of functional excitation and oppression (the latter under ordinary conditions is dynamically stable; disorder of functional equilibrium is conditioned by prevalence of one of syndromes);

- according to functional peculiarities, *YANG* organs (systems) are active, whereas *YIN* organs (systems) – accumulative (tranquility) organs;

- dynamically-stable correlation of *YANG* and *YIN* syndromes, probably correlates with dynamically-stable functional correlation of sympathetic and parasympathetic activity, i.e. with functional-vegetative equilibrium;

- prevalence of *YANG* syndrome over *YIN* syndrome testifies to disorder of vegetative equilibrium with prevalence of sympathetic activity;

- prevalence of *YIN* syndrome over *YANG* syndrome testifies to disorder of vegetative equilibrium with prevalence of parasympathetic activity.

And, the following logical conclusion: Sympathetic and parasympathetic divisions of VNS at organ level act as the executives of functional-informational program of vegetative control. At the same time, the latter is conditioned by vegetative coefficients.

Vegetative coefficients. The developed on the mentioned principles vegetative coefficients (**k**) point to correlation of sympathetic and parasympathetic activity (tab.1). They have been calculated according to the data of examination of 14.304 children and form final functional diagnoses.

Now lets us draw your attention to previously unknown biophysical phenomena that ensure stability of FVD. And we start with the following.

Reality of natural generators of energy is conditioned by two basic positions.

1. Bioelectrical phenomena (processes of distribution and transport of electric charges) are conditioned by the presence in living tissues of a great number of fixed (charged groups of bio macromolecules) and mobile (free electrons and ions) electric charges.

Table 1

ZONE OF ATTENTION OF VEGETATIVE COEFFICIENTS

VALUE of k	ZONE OF FUNCTIONAL ATTENTION	SYMBOL OF ZONE
to 0,75	syndrome of significant para-sympathetic prevalence	PA-s
0,76-0,86	syndrome of significant para-sympathetic prevalence;	PA-e
0,87-0,94	zone of functional compensation of parasympathetic activity	FcP
0,95-1,05	zone of functional-vegetative equilibrium	VE
1,06-1,13	zone of functional compensation of sympathetic activity	FcS
1,14-1,26	syndrome of expressed sympathetic prevalence	SA-e
1,26 and >	syndrome of significant sympathetic prevalence	SA-s

2. Biological systems are natural generators, able to provide transport of free charge carriers into artificially created external circuit.

For realization of biological source of energy, only three components are needed: a) biological system (natural generator of energy); b) chemically inert electrode donor of electrons (DE) – supplier of charge carriers into functional systems through acupunctural zones, and acceptor of electrons (AE) – receiver of free charge carriers through acupunctural zones; c) external circuit with device for control, contacting with electrodes DE and AE.

Source of energy for FVD is the ability of biological systems to generate weak currents into closed circuit. Thus, **diagnostic factor for FVD** is directed transport of free charge carriers (electrons), redistribution of which conditions power-informational transformations. This part requires additional explanation, which we will begin with the question whether there is a relation of functionally active zones (FAZ) of skin to biogenic generation of energy.

Research of their functional activity allowed taking a fresh look at this biophysical phenomenon, and giving answer to the question. They *do* have, and even direct relation! What testifies in favor to this?

1) Movement of charge carriers through biological object (fig.2b) from electrode DE (+) to electrode AE (-), without violation of known direction from (-) to (+) in external circuit. If we add external source of power (battery, fig.6.3a) to the circuit, then the transport of charges becomes dependent on its polarity and it is streaming through the object according to “physical law” from (-) to (+).




Fig.2 Direction of charge carriers through FAZ

2) Movement of charges within closed circuit appears during direct contact of DE and AE with FAZ, and sharply (up to 98,3%) disappears when disconnected. Additionally, transport of charge carriers into external circuit increases up to 42-75% during wet contact of electrodes with FAZ.

3) In the circuit between two single-channel FAZ bioelectrical activity of circuit is in 73,4% higher, than during the contact with FAZ of different functional systems.

Question: do unknown phenomena of FAZ that have direct meaning for vegetative diagnostics exist? Yes, and let us observe them!

Phenomenon of asynchronicity of FAZ or biophysical fractality

We identified five types of temporary symmetrical asynchronicity (fig.3):

- balanced, when during a certain period, bioelectrical activity of two symmetrical FAZ is practically comparable (fig.3.1);




Fig. 3.1

- bilateral, when during a certain period, bioelectrical activity of one of symmetrical FAZ occasionally prevails (fig. 3.2);



Fig. 3.2

- dextral, when during certain period, prevalence of bioelectrical activity of right symmetric FAZ is observed (fig. 3.3);



Fig 3.3

- sinistral, when during certain period, prevalence of bioelectrical activity of left symmetric FAZ is observed (fig. 3.4), and




Fig. 3.4

- mixed, when during a certain period, sequence of previous types of symmetric asynchronicity are met...

At the same time, phenomenon of symmetrical asynchronicity remains also in paired functional systems (fig. 4).

It should be noted, that simultaneous observations

of symmetric asynchronicity in pairs of FAZ of different channels disclosed occasion of different types of interrelations (TS= zone of Test-signal feeding fig.5).






Fig.4 Phenomenon of symmetrical asynchronicity of functional systems.



Signal from right 7ST to 7ST left




Signal from left 7ST to 7ST right




Signal from left 7ST to 6 ST left



Signal from left 6 ST to 7ST left



Signal from left 7ST to 5 ST left



from left 5 ST to 7 ST left

Fig.5. Valvular effect between one-sided FAZ of system ST

Phenomenon of total activity of symmetrical FAZ

Phenomenon of total bioelectrical activity of symmetrical FAZ – is one of basic principles of FVD. It appears in symmetric pairs of FAZ during their simultaneous testing by paired electrode DE with one common output to the controlling device. Thus, for example, triple (during 15-20 minutes) testing of right and left symmetrical FAZ is likely to indicate different values of symmetrical asynchrony. At the same time, if right FAZ will show, for instance, decreasing wave (15-10-5 mcA), then its left analogue, on the contrary, will show rising wave (provisionally opposite values 5-10-15 mcA; fig.6, 6; fig.7; table 2).




Fig.6

Fig.7

Table 2

Individual and total bioelectrical activity of symmetrical FAZ (in mcA)

СТОРОНА SIDE	ІНДИВІДУАЛЬНА І СУМАРНА АКТИВНІСТЬ СИМЕТРИЧНИХ ФАЗ INDIVIDUAL AND TOTAL ACTIVITY OF SYMMETRIC FAZ											
	BL	SP	LI	TE	SI	LU	PC	HT	ST	KI	GB	LR
<i>СПОСТЕРЕЖЕННЯ - SUPERVISION 12.10.2005 (№1)</i>												
ЛІВА - LEFT	1,6	4,4	15	5,0	7,0	0,3	1,0	3,9	2,6	9,0	9,5	1,8
ПРАВА-RIGHT	5,2	3,0	10	5,0	12,0	1,4	12,5	3,0	6,0	11,0	0,8	1,4
Σ	6,8	7,4	24,5	10,0	19,0	1,7	29,5	7,0	8,6	20,0	10,1	3,2
<i>СПОСТЕРЕЖЕННЯ - SUPERVISION 12.10.2005 (№2)</i>												
ЛІВА - LEFT	7,4	10	3,8	2,8	8,5	7,0	6,3	4,0	5,2	2,6	3,5	6,0
ПРАВА-RIGHT	4,2	6,8	6,5	3,3	4,0	7,0	10,5	7,2	2,6	2,0	7,4	3,5
Σ	11,6	16,8	10,3	6,1	12,5	14,0	17,0	11,2	7,8	4,6	11,0	9,5
<i>СПОСТЕРЕЖЕННЯ - SUPERVISION 12.10.2005 (№3)</i>												
ЛІВА - LEFT	1,0	0,9	6,0	0,5	0,8	0,9	0,8	2,5	3,0	3,3	0,8	2,6
ПРАВА-RIGHT	0,7	0,7	6,3	1,5	1,6	0,3	0,5	0,5	1,7	1,6	0,5	2,2
Σ	1,7	1,6	12,0	2,0	2,4	1,2	1,3	3,0	4,7	5,0	1,3	4,8

If two symmetrical FAZs are simultaneously tested with special electrode DE with one common output to gaging unit, we will get stable total results 20-20-20 mcA (fig. 7). At the same time, discrete regularity of the discovered phenomenon should be noted. And, though the latter has no biophysical explanation, its usage in practice of functional-vegetative diagnostics has significant value: for the first time we get stable in time data and twice reduce the number of testings (instead of 24 FAZ, we control total activity of 12 symmetrical pairs).

Other biophysical features will be presented in the second information ...

References

- 1.Val'kova N. YU. Sravnitel'naya otsenka instrumental'noy validnosti anketnogo vegetativnogo testirovaniya i indeksa Kerdo kak metodov integral'noy otsenki vegetativnoy regulyatsii // Problemy nauki i praktiki : regional'nyy podkhod. – Arkhangel'sk, 2000. – S. 118–121.
- 2.Vasilenko A.M. Yegorov V.V., Kozko A.A. Komplementarnaya meditsina na puti k pravovomu regulirovaniyu // Refleksoterapiya i komplementarnaya meditsina. – №1(7), 2014. – S.4-24.
- 3.Vegetativnyye rasstroystva. Klinika, diagnostika, lecheniye /Veyn A. M., Voznesenskaya T. G., Voro-b'yeva O. V. [i dr.] ; pod red. A. Veyna. – M.: Meditsinskoye informatsionnoye agentstvo, 2000. – 750 s. – ISBN 5-89481-066-3
- 4.Vogralik V.G, V. M.V. Punktornaya refleksoterapiya (Chzhen'-tszyu) : monografiya / V. G. Vogralik, M. V. Vogralik. – Gor'kiy, Volgo-Vyatskoye knizhnoye izdatel'stvo, – 1988. – 335 s. – ISBN 5-7420-0073-1

5. Dubrovin Denis Aleksandrovich. Trudnyye voprosy klassicheskoy kitayskoy meditsiny (traktat Nan'tszin) : monografiya / D. A. Dubrovin. – L., Asta Pres, 1991, 223 s. – ISBN 5-85-962-001-02
6. Zhuchenko P.G., Lisov's'ka T.V. Biogal'vanizatsiya, bioenergodiagnostika, bioaktivatsiyina ta antibakterial'na terapiya u virishenni problemi zapal'nikh zakhvoryuvan' v ginekologii. / pid red. d.m.n. V.G.Makatsa. – "UNIVERSUM - Vinnitsya", 1998. – 119 s.
7. Igloukalyvaniye (perevod s v'yetnamskogo). : monografiya / pod obshchey redaktsiyey Khoang Bao Tyao, La Kuang Niyep. – M., Meditsina, 1988. – 672 s. – ISBN 5-225-00299-4
8. Macheret Ye.L., Samosyuk I.Z. Rukovostvo po refleksoterapii : monografiya / Ye. L. Macheret, I. Z. Samosyuk. – K.: "Vishcha shkola", 1982. – 305 s.
9. Netraditsionnyye metody diagnostiki i terapii : monografiya / Samosyuk I.Z., Lysenko V.P., Lima-n'skiy YU.P. [i dr.]; K. : Zdorov'ye, 1994. – 137 s. – ISBN 5-311-00804-0
10. Ovechkin A.M. Osnovy Chzhen'-tszyu terapii : monografiya / A. M. Ovechkin; pod red. prof. V. G. Vogralika. – Saransk, Golos, 1991. – 416 s. – ISBN 5-7585-0007-0
11. Portnov F. G. Elektropunktumaya refleksoterapiya : monografiya / F. G. Portnov. – Riga, Zinatne, 1988. – 352 s. – ISBN 5-7966-0132-6
12. Rozanov A.L. Metod elektropunktumoy diagnostiki "Prognoz" // Refleksoterapiya. – № 1(04)2003. – S. 26-39.
13. Rukovodstvo po refleksoterapii. Elektropunktumaya diagnostika - Nakatani Test : monografiya / Samosyuk I.Z., Fisenko L.I., Chukhrayev N.V. [i dr.]; K. : AO Ukrprofzdravnitsa, 1997 –206 s.
14. Rukovodstvo po refleksoterapii. Elektropunktumaya diagnostika - Nakatani Test : monografiya / Samosyuk I.Z., Fisenko L.I., Chukhrayev N.V. [i dr.]; K. : AO Ukrprofzdravnitsa, 1997. –206 s.
15. Tabeyeva D. M. Rukovodstvo po iglorefleksoterapii : monografiya / D. M. Tabeyeva. – M., Meditsina, 1980. – 560 s.
16. Togoyev A.M., Ponomarev A.V., Usupbekova B.SH., Vasilenko A.M. Vnedreniye metodov reflektornoy diagnostiki i bez lekarstvennoy terapii v otdelenii vosstanovitel'nogo lecheniya gorodskoy polikliniki // Refleksoterapiya. – № 3(14), 2005. – S.50-56.
17. Shen' Chzhitszyan. O tendentsii razvitiya meditsinskoy akupunktury v mire // Refleksoterapiya i komplementarnaya meditsina. №1(1), 2012.–S.14-16.
18. Katsusuke Serizawa/ Effective tsudo therapy : monograph / S. Katsusuke. – Japan publications, INC. – 1984. – 240 s. – ISBN 0-87040-581-0
19. Konig Georg, Wancura Ingrid. Praxis und Theorie der Neuen Chinesischen akupunrtur. Band 1 / G. Konig, I. Wancura. –Wien-Munchen-Bern, 1979, – 402 s. – ISBN 3-85175-294-5 Medicine. China, [1982]. – 408 s. – ISBN 962-06-0287-0
20. Namikoshi Tokujiro/ Shiatsu (Japanese Finger Pressure Therapy) : monograph / T. Namikoshi. – Japan publications, INC. Printed in U.S.A. – 1988. – 82 s. – ISBN 0-87040-169-6
21. Voll Reinhold. Twenty years of electropuncture diagnostic in Germani / R Voll ; Amer. J. Acupuncture. – 1975. – vol. 3. - № 7
22. Wancura Ingrid. Praxis und Theorie der Neuen Chinesischen akupunrtur. Band 2 : monograph / I. Wancura. – Wien-Munchen-Bern, 1983, – 342 s. – ISBN 3-85175-295-3